

COMMENT

Comment to the article: Diaphragmatic patch: A useful adjunct in surgical treatment of recurrent catamenial hemothorax

Comentário ao artigo: Penso diafragmático: um auxiliar útil para o tratamento cirúrgico de hemotórax recorrente catamenial

J.M. Borro

Thoracic Surgery, University Hospital A Coruña, A Coruña, Spain

The present work reviews the catamenial hemothorax and its relation to two potential clinical cases.

Hemothorax is a condition usually associated to thoracic trauma, iatrogenesis, pneumothorax, and malignant or infectious pleural disease. Apart from these clinical entities, the presence of hemothorax is not frequent and its etiology is usually difficult to ascertain. Thoracic endometriosis^{1,2} is one of the rare entities that can be responsible for the presentation of a hemothorax, and the diagnosis is relatively easy when endometriosis foci are found within the thoracic cavity. More frequently, hemothorax is clinically diagnosed in female patients with abdominal endometriosis presenting pneumothorax and/or hemothorax associated with the end phase of the menstrual cycle.^{2,3} Other cases are usually labelled as idiopathic cases.

Regarding the treatment, the use of drainage alone can be sufficient; as a matter of fact, autologous blood was injected in the pleural space as the treatment for pneumothorax since it facilitates the development of pleural adhesences.^{4,5}

Cleaning of pleural cavity through videothoracoscopy, combined with pleural and diaphragmatic abrasion, could facilitate the diagnosis and, at the same time, solve the problem.^{2,3} In case of relapse, total or partial parietal pleurectomy could be a more aggressive option for the management of this entity. In every case, pleural examination and lung palpation should be done in order to identify the presence of lung bullas, endometriosis foci or nodules that should be resected for diagnosis and treatment purposes. The diaphragm should be carefully inspected for the identification and closure with suture of orifices that can be occasionally found.

Talc produces intense adhesences in the pleura, and its use is recommended for the treatment of malignant pleural effusions as palliative treatment. However, talc poudrage

was considered an absolute contraindication to lung transplant. Nowadays, the International Society of Heart and Lung Transplantation still consider talc poudrage as a very high surgical and postoperative risk factor.⁶ All of us who have performed transplant surgery in patients previously treated with talc pleurodesis know the intense pleural reaction against talc, which can be especially problematic when fibrosis affects vascular and mediastinal structures. The use of talc in patients of 30–40 years of age who may require a transplant or any other thoracic surgical intervention in the future should be approached with great caution.

In these cases the reconstruction of the diaphragm with a Gore-Tex patch has been effective,⁷ but it is a very arduous task, expensive and, probably unnecessary, especially when no transdiaphragmatic orifices or significant diaphragmatic pathology are observed.

The end may justify the use of exceptional means in, especially, complex cases, but it never justifies the recommendation of those means as routine treatment.

In conclusion, we should always be extremely thorough in the fulfilment of diagnostic criteria and very cautious when we use or recommend therapeutic procedures that are not yet well established or are not free from future complications.

References

1. El Ghazal R, Fabian T, Ahmed ZA, Moritz ED. Thoracic endometriosis: an unusual cause of hemothorax. *Conn Med.* 2009;73:453–6.
2. Nunes H, Bagan P, Kambouchner M, Martinod E. Thoracic endometriosis. *Rev Mal Respir.* 2007;24:1329–40.
3. Augoulea A, Lambrinouaki I, Christodoulakos G. Thoracic endometriosis syndrome. *Respiration.* 2008;75:113–9.
4. Aihara K, Handa T, Nagai S, Tanizawa K, Watanabe K, Harada Y, et al. Efficacy of blood-patch pleurodesis for secondary spontaneous pneumothorax in interstitial lung disease. *Intern Med.* 2011;50:1157–62.

5. Korasidis S, Andreetti C, D'Andrilli A, Ibrahim M, Ciccone A, Poggi C, et al. Management of residual pleural space and air leaks after major pulmonary resection. *Interact Cardiovasc Thorac Surg*. 2010;10:923–5.
6. Orens JB, Estenne M, Arcasoy S, Conte JV, Corris P, Egan JJ, et al. International Guidelines for the Selection of Lung Transplant Candidates: 2006 Update—a consensus report from the Pulmonary Scientific Council of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. 2006;25:745–55.
7. Nwiloh J. Diaphragmatic patch: A useful adjunct in surgical treatment of recurrent catamenial hemothorax. *Rev Port Pneumol*. 2011;17:278–80.