



PULMONOLOGY

www.journalpulmonology.org


LETTER TO THE EDITOR

Intensity of exercise in people with COPD enrolled in community-based physical activities

Maintaining the benefits of pulmonary rehabilitation (PR) in the long-term is challenging.¹ Community-based physical activity (PA) programmes have emerged as promising strategies to extend these benefits, accommodating different PA facilitators (e.g., supervision, peer support, accessibility, or adequate equipment/infrastructures).²⁻⁴ In addition, community-based PA programmes enable individuals to integrate PA modalities according to their preferences and needs. Nevertheless, community-based PAs should be safe and of at least moderate intensity to align with the PA guidelines.⁵

We examined safety and the intensity level of different community-based PAs in people with COPD who had completed a PR programme.

This observational cross-sectional study, part of two larger trials (PICK UP - NCT04223362, CENTR(AR) - NCT04711057), was approved by the Ethics Committees of *Centro Hospitalar do Baixo Vouga* (Ref.15–05–2019), *Unidade Investigação em Ciências da Saúde – Enfermagem* (Ref.P620–10/2019), and *Administração Regional de Saúde do Centro* (Ref.16/2020). All participants provided informed consent. Privacy policy followed the European Union General Data Protection Regulation 2016/679.

People with stable COPD with a positive risk-benefit analysis (the PA attributes aligned with their characteristics and needs) conducted by the healthcare professionals, who advised their inclusion in community-based PAs, were recruited from five community-based PR programmes. Further details regarding the risk-benefit analysis are provided within the supplementary material. Participants were given a list of available PAs (Table 1 of the supplementary material) and could select those they wanted to try.

Intensity and safety of the community-based PAs were assessed based on data collected from a single exercise session. Heart rate (HR, Polar H10) and percentage of peripheral oxygen saturation (SpO₂, WristOx₂®3150) were continuously monitored during the community-based PAs. Participants' perception of dyspnoea and overall fatigue were assessed every 20-minutes with the modified Borg 0–10 scale. PAs Metabolic Equivalent Tasks' (METs) were estimated using a SenseWear Armband worn by participants during the entire exercise sessions. Due to equipment

constraints SenseWear was not used during pool exercise classes.

Intensity of community-based PAs was determined based on four criteria:

- i. METs: light ≤ 2.9 ; 3 \leq moderate ≤ 5.9 ; vigorous ≥ 6 .⁶
- ii. maximal HR percentage predicted (HRmax_{%predicted}): light ≤ 63 ; 64 \leq moderate ≤ 76 . vigorous ≥ 77 ,⁶ using averaged values collected every 1-second; HRmax_{%predicted} = 220-age.
- iii. iii-iv) dyspnoea and fatigue Borg scores: light ≤ 2 ; 3 \leq moderate ≤ 6 ; vigorous ≥ 7 ,^{6,7} using averaged scores collected every 20 min.

The final PA intensity level was obtained by summing the number of participants categorized within each intensity level (light, moderate and vigorous) according to the four intensity criteria (equally weighted). Safety of community-based PAs was assessed based on SpO₂ and HRmax_{%predicted}. Average and minimum values of SpO₂, and average and maximum values of HRmax_{%predicted} were computed for each participant during PAs. Occurrences of SpO₂ < 88 %, HRmax_{%predicted} > 85 % or any adverse events during the PAs were registered.

Descriptive analyses were conducted. For intensity criteria, absolute frequencies were used and summation presented in bar plots. For safety criteria, mean and 95 % confidence intervals [CI] were displayed. Data analysis was performed using SPSS v.29.0.0.0 and graphs were created using GraphPad Prism 6.

Thirty-two individuals were recruited for this study and 21 agreed to participate. Within these, twelve participants attended senior exercise classes, 10 the gym, and eight pool exercise classes. Participants' characteristics are presented in Table 1.

All community-based PAs were classified as moderate intensity (Fig. 1). During gym, one participant desaturated below 88 %, reaching a minimum of 86 % but recovered within one minute. The 85 % of the HRmax_{%predicted} threshold was surpassed by two participants at the gym (during 58 and 106 s reaching maximum HRmax_{%predicted} of 94 and 89 %, respectively) and by two participants at the senior exercise classes (during 22 and 160 s reaching maximum HRmax_{%predicted} of 88 and 91 %, respectively). No adverse events were registered,

<https://doi.org/10.1016/j.pulmoe.2023.11.001>

2531-0437/© 2023 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1 Participants' characteristics as well as results of intensity and safety criteria per type of community-based physical activity.

	Senior exercise class (n = 12)	Gym (n = 10)	Pool exercise class (n = 8)
Age, years	71.2 ± 8.9	69.6 ± 9.5	69.5 ± 8.8
Sex, male, n (%)	8 (66.7)	10 (100)	6 (75)
BMI, kg/m ²	29.7 ± 6.9	31.4 ± 7.1	32.2 ± 6.8
Smoking status, current/ former/ never, %	0/ 75/ 25	10/ 90/ 0	0/ 75/ 25
Packs/year	29.3 ± 19.6	33.6 ± 24.4	36.8 ± 18.1
FEV ₁ , L/ %predicted	1.4 ± 0.5/ 54.1 ± 12	1.5 ± 0.6/ 55.3 ± 19.3	1.3 ± 0.4/ 52.1 ± 16.5
GOLD grades 1/ 2/ 3/ 4, %	0 / 66.7/ 33.3/ 0	10/ 40/ 40/ 10	12.5/ 50/ 25/ 12.5
GOLD groups A/ B/ E, %	25/ 75/ 0	40/ 60/ 0	37.5/ 62.5/ 0
Charlson Comorbidity Index	4 [3; 5]	4 [3; 5]	4.5 [3; 5]
Mild/ moderate/ severe, %	0/ 66.7/ 33.3	10/ 60/ 30	0/ 50/ 50
Medication, n (%)			
SABA	0	0	0
SAMA	0	0	0
LABA	2 (16.7)	0	0
LAMA	2 (16.7)	2 (20)	0
LAMA + LABA	6 (50)	4 (40)	5 (62.5)
ICS	1 (8.3)	1 (10)	1 (12.5)
ICS + LABA	1 (8.3)	2 (20)	1 (12.5)
ICS + LABA + LAMA	3 (25)	1 (10)	1 (12.5)
LTRA	1 (8.3)	2 (20)	0
Xanthines	0	1 (10)	1 (12.5)
Expectorants	1 (8.3)	3 (30)	0
Oral corticosteroids	0	0	0
Beta blockers	2 (16.7)	1 (10)	1 (12.5)
Intensity criteria			
METs	4.2 [2.5; 4.4]	3.8 [2.8; 4.1]	—
HRmax _{%predicted} (also a safety criterion), %	60.6 ± 9.4	70 ± 8.9	68.5 ± 9.9
Borg fatigue, score	3 [3; 3.8]	3.5 [2.7; 4]	4 [3.3; 4]
Borg dyspnoea, score	3 [2.5; 3.3]	3 [2.7; 3.7]	3 [2.5; 4]
Safety criteria			
HRmax _{%predicted} – maximum, %	74.4 ± 12.5	80.1 ± 8.4	69.8 ± 10.1
SpO ₂ – mean, %	93.5 ± 2.4	92.7 ± 2.7	92.8 ± 2.5
SpO ₂ – minimum, %	93 ± 2.7	91.2 ± 2.9	92.1 ± 5

Values are presented as n (%), mean ± standard deviation or median [interquartile range], unless otherwise stated. Due to technical problems, metabolic equivalent tasks were missing in two participants of the senior exercise classes and in one participant integrating gym.

Legend: BMI: body mass index; FEV₁: forced expiratory volume in one second; GOLD: Global Initiative for chronic obstructive lung disease; HRmax_{%predicted}: maximum HR percentage predicted; ICS: inhaled corticosteroid; LABA: long-acting beta-agonists; LAMA: long-acting muscarinic antagonist; LTRA: leukotriene receptor antagonist; METs – metabolic equivalent tasks; SABA: short-acting beta-agonists; SAMA: short-acting muscarinic antagonist; SpO₂: percentage of peripheral oxygen saturation.

however, due to fatigue one participant had to rest once during gym (two minutes) and twice during the senior exercise class (one and two minutes). Intensity and safety criteria results are presented in [Table 1](#).

Senior, gym and pool exercise classes seem to be safe and of moderate intensity for people with COPD.

Previous studies have found that similar structured senior, gym, or pool exercise classes were safe,^{8,9} with comparable intensity levels reported for aquatic exercises.⁸

Some limitations must be recognized. This was a secondary analysis of larger trials, and no sample size calculation was previously conducted. Our sample included mainly elderly, overweight males with moderate COPD severity and a lower risk of exacerbations. Intensity of PA is influenced by all these individual factors.¹⁰ Additionally, participants were

recruited from community-based PR programmes, thus, our results may not be valid for people with COPD recruited from hospital-based programmes, as the disease severity and complexity are likely to differ.¹¹ Although three relative intensity criteria were used, future studies with larger and more heterogeneous samples are needed. Pooling PA intensity by using different criteria is a strength of this study, as formal guidance on how to assess intensity of community-based PAs is still lacking.¹² To the best of our best knowledge this was the first study assessing intensity of different community-based PA modalities in COPD, in real world settings.

Following PR, based on a careful patient selection, healthcare professionals can now advise people with COPD to integrate gym, senior or pool exercise classes, as these




Fig. 1 Intensity level of community-based physical activities (senior exercise classes [$n = 12$], gym [$n = 10$], and pool exercises classes [$n = 8$]) in people with COPD. Bar graphs represent the sum of the intensity results, i.e., sum of the number of participants categorized within each intensity level (light, moderate and vigorous), according to the four intensity criteria (METs, maximal heart rate percentage predicted [mean values] and dyspnoea and fatigue Borg scores). For each community-based physical activity modality, the total possible sum of intensity results was obtained by multiplying the number of participants by the number of intensity criteria used (senior exercise classes: $12 \times 4 = 48$; gym: $10 \times 4 = 40$; and pool exercise classes: $8 \times 3 = 24$). Due to technical problems, METs were missing in two participants of the senior exercise classes (46 intensity results) and in one participant integrating gym (39 intensity results). Within each coloured rectangle are represented the number of participants categorized according to that intensity criteria in that physical activity intensity level (e.g., within the 12 participants included in the senior exercise classes, according to the Borg dyspnoea criteria [light green], 10 fell within the moderate intensity level and 2 within the light intensity level). Graphs should be read as follows, e.g., in the pool exercise classes, within the 24 intensity results obtained, 18 ($6 + 8 + 4$) fell within the moderate intensity level, 4 ($2 + 2$) within the light intensity level and 2 within the vigorous intensity level.

Legend: HRmax %predicted: maximum HR percentage predicted; METs – metabolic equivalent tasks.

seem to be safe, of moderate intensity, and may enhance long-term maintenance of PR benefits.

CRedit authorship contribution statement

P.R., D.B. and A.M. - conceptualization; P.R., A.T., R.P. and C.S. - data curation; P.R., D.B. and A.M. - formal analyses, investigation and methodology; AM and PR – funding acquisition; AM – Project administration; AM and DB – Supervision; PR – Writing - original draft; P.R., A.T., R.P., C.S., DB and AM – Writing review and editing.

Data availability statement

Data are available upon reasonable request to the corresponding author.

Declaration of Competing Interest

The authors have no conflicts of interest.

Funding

This work was funded by the project “CENTR(AR): pulmões em andamento” by Programa de Parcerias para o Impacto, Portugal Inovação Social through Programa Operacional Inclusão Social e Emprego (POISE-03-4639-FSE-000597); by Fundação para a Ciência e a Tecnologia (SFRH/BD/148738/2019 and COVID/BD/153476/2023) by Fundo Social Europeu

through Programa Operacional Regional Centro, and by Programa Operacional Competitividade e Internacionalização (COMPETE 2020 - POCI-01-0145-FEDER-007628; UIDB/04501/2020 and UIDP/04501/2020).

Acknowledgements

The authors would like to thank Guilherme Rodrigues, Cíntia Dias, Ana Sofia Grave, Sara Almeida, Eduardo Samuel Santos, Inês Agostinho and Carolina Monteiro for their contribution during data collection.

Supplementary materials

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.pulmoe.2023.11.001](https://doi.org/10.1016/j.pulmoe.2023.11.001).

References

1. Spruit MA, Singh SJ, Garvey C, et al. An official american thoracic society/european respiratory society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med.* 2013;188(8):e13–64.
2. Meshe OF, Bungay H, Claydon LS. Participants' experiences of the benefits, barriers and facilitators of attending a community-based exercise programme for people with chronic obstructive pulmonary disease. *Health Soc Care Community.* 2020;28(3):969–78.
3. Desveaux L, Harrison S, Lee A, et al. "We are all there for the same purpose": support for an integrated community exercise

- program for older adults with HF and COPD. *Heart Lung*. 2017;46(4):308–12.
4. Robinson H, Williams V, Curtis F, et al. Facilitators and barriers to physical activity following pulmonary rehabilitation in COPD: a systematic review of qualitative studies. *NPJ Prim Care Respir Med*. 2018;28(1):19.
 5. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451–62.
 6. American College of Sports Medicine. *ACSM's Guidelines for Exercise Testing and Prescription*. 10th ed. Wolters Kluwer; 2018.
 7. Borg G. Borg's perceived exertion and pain scales: human kinetics; 1998.
 8. McNamara RJ, McKeough ZJ, McKenzie DK, et al. Water-based exercise training for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2013(12):Cd008290.
 9. Butler SJ, Desveaux L, Lee AL, et al. Randomized controlled trial of community-based, post-rehabilitation exercise in COPD. *Respir Med*. 2020;174:106195.
 10. Farooqi N, Carlsson M, Häglin L, et al. Energy expenditure in women and men with COPD. *Clin Nutr ESPEN*. 2018;28:171–8.
 11. Spruit MA, Wouters EFM. Organizational aspects of pulmonary rehabilitation in chronic respiratory diseases. *Respirology*. 2019;24(9):838–43.
 12. Rebelo P, Brooks D, Marques A. Measuring intensity during free-living physical activities in people with chronic obstructive pulmonary disease: a systematic literature review. *Ann Phys Rehabil Med*. 2022;65(2):101607.
- P. Rebelo^{a,b}, A. Teixeira^c, R. Pinto^c, C. Santos^c, D. Brooks^{d,e}, A. Marques^{a,b,*}
- ^a *Lab3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences, University of Aveiro, Aveiro, Portugal*
- ^b *iBiMED - Institute of Biomedicine, Department of Medical Sciences, University of Aveiro, Aveiro, Portugal*
- ^c *City council of Estarreja, Sports division, Estarreja, Portugal*
- ^d *School of Rehabilitation Science, McMaster University, Hamilton, ON, Canada*
- ^e *West Park Healthcare Centre, Toronto, ON, Canada*
- * Corresponding author:
E-mail address: amarques@ua.pt (A. Marques).
Received 11 July 2023; Accepted 2 November 2023
Available online xxx