

2. Chung F, Subramanyam R, Liao P, Sasaki E, Shapiro C, Sun Y. High STOP-Bang score indicates a high probability of obstructive sleep apnoea. *Br J Anaesth*. 2012;108:768–75.
3. Liao P, Yegneswaran B, Vairavanathan S, Zilberman P, Chung F. Postoperative complications in patients with obstructive sleep apnea: a retrospective matched cohort study. *Can J Anaesth*. 2009;56:819–28.
4. Gupta RM, Parvizi J, Hanssen AD, Gay PC. Postoperative complications in patients with obstructive sleep apnea syndrome undergoing hip or knee replacement: a case-control study. *Mayo Clin Proc*. 2001;76:897–905.
5. Memtsoudis S, Liu SS, Ma Y, Chiu YL, Walz JM, Gaber-Baylis LK, Mazumdar M. Perioperative pulmonary outcomes in patients with sleep apnea after noncardiac surgery. *Anesth Analg*. 2011;112:113–21.

Helder Pereira^a, Daniela Xará^a, Júlia Mendonça^a, Alice Santos^a, Fernando Abelha^{a,b,*}

^a Department of Anesthesiology, Centro Hospitalar de São João, Porto, Portugal

^b Anesthesiology and Perioperative Care Unit, Surgical Department of Medical School of Porto, Porto, Portugal

* Corresponding author.

E-mail address: fernando.abelha@gmail.com (F. Abelha).

Video-mediastinoscopy is still the gold standard



A video-mediastinoscopia é ainda o gold-standard

Dear Editor,

We read with great interest the article by Bugalho et al., entitled “Endobronchial ultrasound-guided transbronchial needle aspiration for lung cancer diagnosis and staging in 179 patients”¹ as well as the editorial by Herth entitled “Access to the mediastinum—The standard has changed”.²

In fact, for patients with lung cancer, despite improvements in the accuracy of imaging modalities over the last decade, invasive mediastinal lymph node staging remains necessary in cases of mediastinal lymph node enlargement, positron emission tomography (PET) positive mediastinal and/or hilar lymph nodes and/or a centrally located tumor.³ For a long time, cervical mediastinoscopy has been considered the gold standard in mediastinal staging, given the high negative predictive value (NPV) if well performed. However, during the last decade, oesophageal ultrasound-guided fine needle aspiration (EUS-FNA) followed by endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) has emerged as a minimally invasive alternative, reducing the need for a cervical mediastinoscopy as a first-line staging procedure. Nevertheless, when the result of endoscopic staging appears negative, a subsequent mediastinoscopy is currently recommended to exclude mediastinal lymph node metastases in patients with clinical suspicion.² But, since the sensitivity of EBUS-FNA seems to exceed that of mediastinoscopy,⁴ there is a tendency to cut down on the need for surgical confirmation. Accordingly, in routine practice an additional mediastinoscopy is often regarded as overdone. However, only recently the combination of endosonography followed by mediastinoscopy was shown to be more accurate in mediastinal nodal staging than just mediastinoscopy alone. In patients with non-small-cell lung cancer and an indication for mediastinal staging, performing a cervical mediastinoscopy after a negative result of endosonography reduced the number of futile thoraco-

tomies by 50%. Overall, an average of 8.8 patients had to undergo an additional mediastinoscopy to find one false-negative result of endosonography, but when only patients with suspicious mediastinal lymph nodes on FDG-PET are taken into account, this NNT comes down to 6.1 patients.⁵

We agree with Dr. Herth. The standard has changed. In fact, EBUS and EUS should be used in conjunction with PET and considered as an alternative to mediastinoscopy. However, the gold standard is still video-mediastinoscopy, because in patients with a high probability of mediastinal metastases, a cervical mediastinoscopy should not be omitted after a negative result of endosonography, not even when the aspirate seems representative, based on the presence of a sufficient number and maturation of lymphocytes.

References

1. Bugalho A, Ferreira D, Barata R, Rodrigues C, Dias SS, Medeiros F, et al. Endobronchial ultrasound-guided transbronchial needle aspiration for lung cancer diagnosis and staging in 179 patients. *Rev Port Pneumol*. 2013;19:192–9.
2. Herth FJ. Access to the mediastinum—the standard has changed. *Rev Port Pneumol*. 2013;19:190–1.
3. De Leyn P, Lardinois D, van Schil PE, Rami-Porta R, Passlick B, Zielinski M, et al. ESTS guidelines for preoperative lymph node staging for non-small cell lung cancer. *Eur J Cardiothorac Surg*. 2007;32:1–8.
4. Gu P, Zhao YZ, Jiang LY, Zhang W, Xin Y, Han BH. Endobronchial ultrasound-guided transbronchial needle aspiration for staging of lung cancer: a systematic review and meta-analysis. *Eur J Cancer*. 2009;45:1389–96.
5. Verhagen AF, Schuurbiens OC, Looijen-Salamon MG, van der Heide SM, van Swieten HA, van der Heijden EH. Mediastinal staging in daily practice: endosonography, followed by cervical mediastinoscopy. Do we really need both? *Interact CardioVasc Thorac Surg*. 2013;17:823–8.

M. Guerra

Serviço de Cirurgia Cardiorácica, Centro Hospitalar de Vila Nova de Gaia/Espinho, EPE, Vila Nova de Gaia, Portugal

E-mail address: miguel david guerra@yahoo.com